

Elucidating the Vacuum Structure of the Aoki Phase

V. Azcoitiⁱ, G. Di Carloⁱⁱ, E. Follanaⁱ and A. Vaqueroⁱⁱⁱ.

October 5, 2012

Abstract

In this paper, we discuss the vacuum structure of QCD with two flavors of Wilson fermions, inside the Aoki phase. We provide numerical evidence, coming from HMC simulations in 4^4 , 6^4 and 8^4 lattices, supporting a vacuum structure for this model at strong coupling more complex than the one assumed in the standard wisdom, with new vacua where the expectation value of $i\bar{\psi}\gamma_5\psi$ can take non-zero values, and which can not be connected with the Aoki vacua by parity-flavour symmetry transformations.

1 Introduction

Two and three colors QCD with unimproved Wilson fermions started to be simulated in the early 80's [1, 2]. The complexity of the phase structure of this model was known from the very beginning, and indeed the existence of a phase at finite lattice spacing, a , with spontaneous parity and flavor symmetry breaking, was conjectured for this model by Aoki in the middle 80's [3, 4]. From that time on, much work has been done in order to confirm this pattern of spontaneous symmetry breaking, and to establish a quantitative phase diagram for lattice QCD with Wilson fermions. References [5–26] are an incomplete list of the work done along these years.

Aoki's conjecture has been supported not only by numerical simulations but also by theoretical studies based on the linear sigma model [6] and on applying chiral perturbation theory (χPT) to the continuum effective lagrangian [11]. The latter analysis predicts, near the continuum limit, two possible scenarios, depending on the sign of an unknown low-energy coefficient. In one scenario, flavor and parity are spontaneously broken, and there is an Aoki phase with a nonzero value only for the $i\bar{\psi}\gamma_5\tau_3\psi$ condensate. In the other one (the first-order scenario) there is no spontaneous symmetry breaking.

ⁱDepartamento de Física Teórica, Facultad de Ciencias, Universidad de Zaragoza, Cl. Pedro Cerbuna 12, E-50009 Zaragoza (Spain)

ⁱⁱINFN, Laboratori Nazionali del Gran Sasso, I-67100 Assergi, L'Aquila (Italy)

ⁱⁱⁱComputation-based Science and Technology Research Center (CaSToRC), The Cyprus Institute, 20 Constantinou Kavafi Street, Nicosia 2121, (Cyprus)

This standard picture for the Aoki phase was questioned three years ago by three of us in [21], where we conjectured on the appearance of new vacua in the Aoki phase, which can be characterized by a non-vanishing vacuum expectation value of the flavor singlet pseudoscalar condensate $i\bar{\psi}\gamma_5\psi$, and which can not be connected with the Aoki vacua by parity-flavor symmetry transformations. However Sharpe pointed out in [22] that our results, based on an analysis of the Probability Distribution Function (PDF) of fermion bilinears, could still be reconciled with the standard scenario, and that otherwise we would question also the validity of the χPT analysis.

For the last few years we have performed an extensive research on the vacuum structure of the Aoki phase [23–25], in order to find out if our alternative vacuum structure, derived from the PDF analysis, was realized or not. The purpose of this paper is to clarify these issues by reporting the results of our investigations which, as will be shown along this article, provide evidence of a more complex vacuum structure in the Aoki phase of two-flavor QCD, as conjectured in [21].

The outline of the paper is as follows. Section 2 summarizes the standard picture on the phase diagram of QCD with two flavors of unimproved Wilson fermions and our alternative scenario, derives some interesting expressions to be compared with continuum results, and analyzes the relevance of a discrete symmetry P' , composition of parity and discrete flavor transformations, which was introduced by Sharpe and Singleton [11] as an attempt to reconcile the Vafa-Witten theorem on the impossibility to break spontaneously parity in a vector-like theory, with the existence of the Aoki phase. Section 3 contains technical data of our HMC simulations of QCD with two flavors of Wilson fermions, inside and outside the Aoki phase, performed without external sources and also with a twisted mass term in the action. The results of our analysis of the PDF of the operator Q , an order parameter for the P' symmetry, are reported in Section 4. This Section contains what is, in our opinion, the strongest indication on the existence of a vacuum structure in the Aoki phase more complex than the standard one. In Section 5 we show how we were able to perform a direct measurement of $\langle (i\bar{\psi}\gamma_5\psi)^2 \rangle$ in the Aoki phase, which produced a non-zero value for this operator in the 6^4 lattice, of the same order of magnitude as $\langle (i\bar{\psi}_u\gamma_5\psi_u)^2 \rangle$ and $\langle (i\bar{\psi}\gamma_5\tau_3\psi)^2 \rangle$ (both are non-vanishing in the Aoki phase in the standard picture). Finally Section 6 contains our conclusions.

2 The Aoki phase for the two flavor model

The fermionic Euclidean action of QCD with two degenerate Wilson flavors is

$$S_F = \bar{\psi}_u W(\kappa) \psi_u + \bar{\psi}_d W(\kappa) \psi_d, \quad (1)$$

where $W(\kappa)$ is the Dirac-Wilson operator, and κ is the hopping parameter, related to the adimensional bare fermion mass m_0 by $\kappa = 1/(8 + 2m_0)$. The standard wisdom on the phase diagram of this model in the gauge coupling β , κ plane is the

one shown in figure 1. The two different regions observed in this phase diagram, A and B, can be characterized as follows: in region A parity and flavor symmetries are realized in the vacuum, which is supposed to be unique. The continuum limit should be obtained by taking the $g^2 \rightarrow 0$, $\kappa \rightarrow 1/8$ limit from within region A, which we will call the QCD region. In region B, on the contrary, parity and flavor symmetries are spontaneously broken, there are many degenerate vacua connected by parity-flavor transformations in this region, and the Gibbs state is therefore made up from many degenerate vacuum states. In what follows we will call region B the Aoki region.




Figure 1: Aoki (B) and physical (A) region in the (β, κ) plane. Adapted from [17] by courtesy of the authors.

2.1 Standard Wisdom

The standard order parameters to distinguish the Aoki region from the QCD region are $i\bar{\psi}\gamma_5\tau_j\psi$ and $i\bar{\psi}\gamma_5\psi$, with τ_j the three Pauli matrices. For the standard election, $j = 3$, they can be written as a function of the up and down quark fields as follows

$$\begin{aligned} i\bar{\psi}\gamma_5\tau_3\psi &= i\bar{\psi}_u\gamma_5\psi_u - i\bar{\psi}_d\gamma_5\psi_d, \\ i\bar{\psi}\gamma_5\psi &= i\bar{\psi}_u\gamma_5\psi_u + i\bar{\psi}_d\gamma_5\psi_d. \end{aligned} \tag{2}$$

The standard wisdom for the Aoki phase can be summarized by the following two equations [4]

$$\begin{aligned}\langle i\bar{\psi}\gamma_5\tau_3\psi \rangle &\neq 0, \\ \langle i\bar{\psi}\gamma_5\psi \rangle &= 0.\end{aligned}\tag{3}$$

These equations should hold in the vacuum selected when we add a twisted mass term to the euclidean Wilson action (1),

$$S_{m_t} = m_t i\bar{\psi}\gamma_5\tau_3\psi,\tag{4}$$

which breaks both parity and flavor symmetries, and take the vanishing twisted mass limit once the infinite degrees of freedom limit has been performed. The first of the two condensates $i\bar{\psi}\gamma_5\tau_3\psi$ breaks both parity and flavor symmetries. The non-vanishing vacuum expectation value of this condensate signals the spontaneous breaking of the $SU(2)$ flavor symmetry down to $U(1)$, with two Goldstone charged pions. Notwithstanding parity is spontaneously broken if the first equation in (3) holds, the vacuum expectation value of the flavor diagonal condensate $\langle i\bar{\psi}\gamma_5\psi \rangle$ vanishes. Indeed it is also an order parameter for a discrete symmetry P' , composition of parity and discrete flavor rotations, which is assumed to be realized [11].

The Aoki phase can then be characterized by the presence of two massless charged pions and a massive neutral pion which becomes massless just at the critical line separating this phase from the physical one. In addition the η -meson, the mass of which can be extracted from the long distance behavior of the two-point correlation function of the flavor singlet pseudoscalar operator $i\bar{\psi}\gamma_5\psi$, stays massive even at the critical line as a consequence of the $U(1)_A$ axial anomaly.

2.2 Our alternative Scenario

Three years ago this standard scenario was revised in [21] with the help of the probability distribution function (PDF) of the parity-flavor fermion bilinear order parameters [27][28]. We found that if the spectral density $\rho_U(\lambda, \kappa)$ of the overlap hamiltonian $\gamma_5 W(\kappa)$, or Hermitian Dirac-Wilson operator, in a fixed background gauge field U were not symmetric in λ in the thermodynamic limit for the relevant gauge configurations, Hermiticity of $i\bar{\psi}_u\gamma_5\psi_u + i\bar{\psi}_d\gamma_5\psi_d$ would be violated at finite β . Assuming that the Aoki phase ends at finite β , this would imply the loss of any physical interpretation of this phase in terms of particle excitations. If on the contrary the spectral density $\rho_U(\lambda, \kappa)$ of the Hermitian Dirac-Wilson operator in a fixed background gauge field U is symmetric in λ , we argued in [21] on the appearance of other phases, which can be characterized by a non-vanishing vacuum expectation value of $i\bar{\psi}_u\gamma_5\psi_u + i\bar{\psi}_d\gamma_5\psi_d$, and which can not be connected with the Aoki vacua by parity-flavor symmetry transformations. We summarize here the main steps of [21].

The key point was to study the Fourier transform of the PDF of the fermion bilinear order parameters, $P(q)$, defined as

$$P(q) = \frac{\int [dU] [d\bar{\psi} d\psi] \exp\{-S_{YM} + \bar{\psi} W(\kappa) \psi + \frac{iq}{V} \int d^4x O(x)\}}{\int [dU] [d\bar{\psi} d\psi] \exp\{-S_{YM} + \bar{\psi} W(\kappa) \psi\}}, \quad (5)$$

where O is any fermion bilinear operator

$$O(x) = \bar{\psi}(x) \tilde{O} \psi(x), \quad (6)$$

with \tilde{O} a matrix with Dirac, color and flavor indices, and V is the number of lattice points. Integrating out the fermion fields one gets

$$P(q) = \frac{\int [dU] e^{-S_{YM}} \det(W(\kappa) + \frac{iq}{V} \tilde{O})}{\int [dU] e^{-S_{YM}} \det W(\kappa)}, \quad (7)$$

which can also be expressed as the following mean value

$$P(q) = \left\langle \frac{\det(W(\kappa) + \frac{iq}{V} \tilde{O})}{\det W(\kappa)} \right\rangle, \quad (8)$$

computed in the effective gauge theory with the integration measure

$$[dU] e^{-S_{YM}} \det W(\kappa).$$

The q -derivatives of $P(q)$ give us the moments of the PDF of our fermion bilinear

$$\left. \frac{d^n P(q)}{dq^n} \right|_{q=0} = i^n \left\langle \left(\frac{1}{V} \int d^4x O(x) \right)^n \right\rangle. \quad (9)$$

For the case we are interested in, and if we call $P_j(q), P_0(q)$ the PDF of

$$\begin{aligned} c_j &= i \bar{\psi} \gamma_5 \tau_j \psi, \\ c_0 &= i \bar{\psi} \gamma_5 \psi. \end{aligned} \quad (10)$$

In momentum space, we have

$$\begin{aligned} P_1(q) &= P_2(q) = P_3(q) = \left\langle \prod_j \left(-\frac{q^2}{V^2 \lambda_j^2} + 1 \right) \right\rangle, \\ P_0(q) &= \left\langle \prod_j \left(\frac{q}{V \lambda_j} + 1 \right)^2 \right\rangle, \end{aligned} \quad (11)$$

where λ_j are the real eigenvalues of the one flavor Hermitian Dirac-Wilson operator $\bar{W}(\kappa) = \gamma_5 W(\kappa)$.

Since c_0, c_j are order parameters for the symmetries of the lattice action and we are not selecting a particular vacuum state, their first moment will vanish always, independently of the realization of the symmetries. The first non-vanishing moment, if the symmetry is spontaneously broken, will be the second:

$$\begin{aligned}
\langle c_0^2 \rangle &= 2 \left\langle \frac{1}{V^2} \sum_j \frac{1}{\lambda_j^2} \right\rangle - 4 \left\langle \left(\frac{1}{V} \sum_j \frac{1}{\lambda_j} \right)^2 \right\rangle, \\
\langle c_3^2 \rangle &= 2 \left\langle \frac{1}{V^2} \sum_j \frac{1}{\lambda_j^2} \right\rangle.
\end{aligned} \tag{12}$$

In the QCD region flavor symmetry is realized. The PDF of c_3 will be then $\delta(c_3)$ and $\langle c_3^2 \rangle = 0$. We get then

$$\langle c_0^2 \rangle = -4 \left\langle \left(\frac{1}{V} \sum_j \frac{1}{\lambda_j} \right)^2 \right\rangle,$$

which should vanish since parity is also realized in this region. Furthermore a negative value of $\langle c_0^2 \rangle$ would violate Hermiticity of $i\bar{\psi}\gamma_5\psi$.

In the Aoki region [4] there are vacuum states in which the condensate c_3 (10) takes a non-vanishing vacuum expectation value. This implies that the PDF $P(c_3)$ will not be $\delta(c_3)$ and therefore $\langle c_3^2 \rangle$ (12) will not vanish. Indeed expression (12) for $\langle c_3^2 \rangle$ seems to be consistent with the Banks and Casher formula [29] which relates the spectral density of the Hermitian Dirac-Wilson operator at the origin with the vacuum expectation value of c_3 [11].

If, on the other hand, $\langle i\bar{\psi}\gamma_5\psi \rangle = 0$ in one of the Aoki vacua, as conjectured in [4], $\langle i\bar{\psi}\gamma_5\psi \rangle = 0$ in all the other vacua that are connected with the standard Aoki vacuum by a parity-flavor transformation, since $i\bar{\psi}\gamma_5\psi$ is invariant under flavor transformations and changes sign under parity. Therefore if we assume that these are all the degenerate vacua, we should conclude that $P(c_0) = \delta(c_0)$ and $\langle c_0^n \rangle = 0$, which would imply an infinite tower of non trivial relations, one for each even moment of the PDF. We write here the simplest one,

$$\left\langle \frac{1}{V^2} \sum_j \frac{1}{\lambda_j^2} \right\rangle = 2 \left\langle \left(\frac{1}{V} \sum_j \frac{1}{\lambda_j} \right)^2 \right\rangle \neq 0. \tag{13}$$

A possible scenario which was discussed in [21] is the one that emerges if we assume a symmetric distribution for the eigenvalues of the hermitian Wilson operator. As discussed in [21] this assumption is necessary in order to preserve hermiticity. If we take this assumption in the most naive way we get

$$\langle c_3^2 \rangle - \langle c_0^2 \rangle = 4 \left\langle \left(\frac{1}{V} \sum_j \frac{1}{\lambda_j} \right)^2 \right\rangle = 0, \tag{14}$$

and therefore we should conclude that new vacua characterized by a non vanishing vacuum expectation value of the singlet-flavor pseudoscalar operator $i\bar{\psi}\gamma_5\psi$ must exist besides the Aoki vacua. Nevertheless as Sharpe noticed in [22], sub-leading contributions to the spectral density may affect equation (14) in the ε -regime in

such a way that, not only $\langle c_0^2 \rangle$, but every even moment of $i\bar{\psi}\gamma_5\psi$ would vanish, restoring the standard Aoki picture.

As we will show in this article equation (14) turns out not to be realistic in the Aoki phase, as follows from numerical simulations. However the thesis of Sharpe, although possible and inspired in the absence of our conjectured new vacua in the chiral effective lagrangian approach, would enforce as discussed before an infinite series of *sum rules*, similar to those found by Leutwyler and Smilga in the continuum [30]. The main purpose of this paper is to clarify these issues.

2.3 Some interesting relations

As previously discussed flavor and parity symmetries should be realized in the vacuum of the physical phase. This phase can therefore be characterized, in what concerns its low energy spectrum, by the existence of three degenerate massive pions, which become massless at the critical line, and the η -meson, which is massive even at the critical line because of the axial anomaly. When we cross the critical line and enter into the Aoki phase, the neutral pion becomes massive whereas the charged pions are massless since they are the two Goldstone bosons associated to the spontaneous breaking of the $SU(2)$ flavor group to a $U(1)$ group.

The susceptibilities of the neutral pion χ_{π^0} and the eta-meson χ_η in the physical phase are the corresponding integrated two-point correlation functions

$$\chi_{\pi^0} = \sum_x \langle i\bar{\psi}\gamma_5\tau_3\psi(x) i\bar{\psi}\gamma_5\tau_3\psi(0) \rangle = \frac{2}{V} \left\langle \sum_{i=1}^V \frac{1}{\lambda_i^2} \right\rangle, \quad (15)$$

$$\chi_\eta = \sum_x \langle i\bar{\psi}\gamma_5\psi(x) i\bar{\psi}\gamma_5\psi(0) \rangle^1 = \frac{2}{V} \left\langle \sum_{i=1}^V \frac{1}{\lambda_i^2} \right\rangle - \frac{4}{V} \left\langle \left[\sum_{i=1}^V \frac{1}{\lambda_i} \right]^2 \right\rangle. \quad (16)$$

The rightmost hand sides of these equations are just the second moments of the PDF of $i\bar{\psi}\gamma_5\tau_3\psi$ and $i\bar{\psi}\gamma_5\psi$ (see equations (12)) multiplied by the corresponding volume factor. Therefore equations (15), and (16) give us the following relation between χ_η , χ_{π^0} and the trace of the inverse one-flavor hermitian Dirac-Wilson operator:

$$\chi_\eta = \chi_{\pi^0} - \frac{4}{V} \langle \text{Tr}^2 (\gamma_5 W^{-1}(\kappa)) \rangle, \quad (17)$$

to be compared with the well known relation between the eta, pion and topological susceptibilities

$$\chi_\eta = \chi_{\pi^0} + 4 \frac{\chi_T}{m^2}, \quad (18)$$

which holds in the continuum and also in the Ginsparg-Wilson regularization at finite lattice spacing.

¹There are no disconnected contributions in this phase.

The first interesting observation that follows from equations (17), and (18) is the suggestive relation between the topological susceptibility χ_T and the trace of the inverse hermitian Dirac-Wilson operator,

$$\frac{\chi_T}{m^2} = -\frac{1}{V} \langle \text{Tr}^2 (\gamma_5 W^{-1}(\kappa)) \rangle, \quad (19)$$

which should hold near the continuum limit. This suggests also the following relation between the trace of the inverse hermitian Dirac-Wilson operator, quark mass m , and the density of topological charge $q = Q/V$ [31, 32]

$$\frac{q}{m} = \frac{1}{V} \text{Tr} (\gamma_5 W^{-1}(\kappa)), \quad (20)$$

A second interesting observation that follows from (17) concerns the behavior of

$$\frac{4}{V} \langle \text{Tr}^2 (\gamma_5 W^{-1}(\kappa)) \rangle = \frac{4}{V} \left\langle \left[\sum_{i=1}^V \frac{1}{\lambda_i} \right]^2 \right\rangle \quad (21)$$

when we approach the critical line from the physical phase. Indeed in the physical phase this quantity should be finite because the pion and eta susceptibilities are finite. However when approaching the critical line it should diverge in order to compensate the divergence of the pion susceptibility keeping χ_η finite in (17), in deep similarity with the divergence of the topological susceptibility divided by the square quark mass in the continuum and Ginsparg-Wilson regularization in the chiral limit. The origin of this divergence lies in the accumulation of eigenvalues of the hermitian Dirac-Wilson operator near the origin.

The divergence of (21) at the critical line is slower than V in such a way as to keep a parity-flavor symmetric vacuum, as corresponds to a second order phase transition. On the other hand if the standard scenario for the Aoki phase is realized, the higher accumulation of eigenvalues of the hermitian Dirac-Wilson operator at the origin in this phase should be enough to give a finite contribution to $\langle c_i^2 \rangle$ (second equation in (12)). Indeed if the Aoki vacuum selected with a twisted mass term plus those obtained by parity-flavor symmetry transformations are the only degenerate vacua in this phase, the following relation should hold [22, 27]

$$\langle c_3^2 \rangle = \frac{\Sigma^2}{3}, \quad (22)$$

where Σ is the mean value of c_3 in the vacuum selected by the twisted mass term. In Section 5 we will report the results of a check of equation (22).

The standard scenario for the Aoki phase requires also that the singlet flavor pseudoscalar operator $i\bar{\psi}\gamma_5\psi$ takes a vanishing vacuum expectation value. Taking into account the first equation in (12), this requirement implies that $\frac{4}{V} \langle \text{Tr}^2 (\gamma_5 W^{-1}(\kappa)) \rangle$ in (21) should diverge as V , to compensate the first contribution to $\langle c_0^2 \rangle$ in (12).

2.4 Spontaneous Symmetry Breaking of P' ?

Years ago Sharpe and Singleton suggested [11] that a symmetry P' , composition of parity and flavor transformations, associated to a discrete subgroup of the parity-flavor continuous group, and acting on pure gluonic operators as parity, could still be realized in the Aoki phase, notwithstanding parity and flavor are spontaneously broken in this phase. Their motivation for such a proposal was the attempt to reconcile the Vafa-Witten theorem [33] on the impossibility to break spontaneously parity in a vector-like theory, with the existence of the Aoki phase. Their point was that Vafa-Witten theorem, although it does not apply to fermionic order parameters, could still apply to pure gluonic operators. Then if the realization of P' does not require a vanishing Aoki condensate, and P' acts on pure gluonic operators as parity, the existence of the Aoki phase besides the realization of P' in the vacuum would not be in conflict with the Vafa-Witten theorem for pure gluonic operators. Furthermore the realization of P' would be quite useful since it would give a simple explanation for the tower of sum rules (13) required in the standard scenario. Indeed the flavor-singlet pseudoscalar condensate $i\bar{\psi}\gamma_5\psi$ is an order parameter for P' .

Although the main motivation for the introduction of the P' symmetry, the realization of Vafa-Witten theorem for pure gluonic operators, has become less relevant on the light of later works on this subject [34]-[37], the implications on the standard scenario for the Aoki phase of the realization of this symmetry are still relevant.

A possible election for P' , the realization of which would not be in conflict with non vanishing condensates $\langle i\bar{\psi}\gamma_5\tau_j\psi \rangle$ for $j = 1, 2, 3$ is the composition of parity with the Z_4 subgroup of the $SU(2)$ flavor group generated by

$$\frac{i}{\sqrt{3}}(\tau_1 + \tau_2 + \tau_3).$$

In this context let us consider the operator

$$Q = \frac{1}{V} \text{Tr}(\gamma_5 W^{-1}(\kappa)) = \frac{1}{V} \sum_{j=1}^V \frac{1}{\lambda_j}, \quad (23)$$

which is a non local order parameter for P' . This operator, although non-local, is not singular since exact zero modes have vanishing integration measure in the Wilson formulation. We will assume in what follows that Q is an intensive operator. Notice that any local operator should be intensive but any intensive operator is not necessarily local. Our assumption is based in the following argument: $\langle Q \rangle = \frac{1}{2V} \langle \sum_x \bar{\psi}(x)\gamma_5\psi(x) \rangle$, $\bar{\psi}(x)\gamma_5\psi(x)$ being a local fermionic operator. Our prejudices tell us that $\frac{1}{2V} \sum_x \bar{\psi}(x)\gamma_5\psi(x)$, as any intensive operator, should be self-averaging. In other words, if the system size is very large, a single thermalized gauge configuration should be enough to get $\frac{1}{2V} \langle \sum_x \bar{\psi}(x)\gamma_5\psi(x) \rangle$, and this seems indeed to be the case in the physical phase, where the PDF of Q approaches a delta, as shown in Section 4. Furthermore equations (17) and (18) strongly suggest that,

near the continuum limit, Q would be essentially the density of topological charge, which indeed is an intensive operator.

Hence were the P' symmetry realized in the vacuum, the PDF of (23) would be, according to our assumption, a $\delta(x)$ distribution. But the second moment of this PDF,

$$\frac{1}{V^2} \left\langle \left[\sum_{i=1}^V \frac{1}{\lambda_i} \right]^2 \right\rangle,$$

needs to be finite in the standard scenario for the Aoki phase in order to realize the first sum rule (13). Therefore the standard scenario requires the spontaneous symmetry breaking of P' , but then there are no symmetry reasons to justify this tower of sum rules. In Section 4 we will report numerical results for the PDF of (23), both in the Aoki and in the physical phase.

3 The simulations

In order to determine the phase structure and the properties of the vacuum, we need to measure the PDF 's of the parity-flavor order parameters described in the previous sections, including the Q operator (23). So we decided to carry out HMC simulations of QCD with two flavors of Wilson fermions, inside and outside the Aoki phase, and mainly without external sources, although some simulations with a twisted mass term in the action have also been performed.

We remove the external sources because of several facts:

1. The PDF formalism requires the removal of any external sources in the action. We just perform one long simulation in the Gibbs state.
2. The addition of a twisted mass external source, as has been done in past simulations of the Aoki phase, automatically selects the vacuum where the standard properties of the Aoki phase are verified [21]. This point could explain why no one ever saw a new Aoki phase like the one we are proposing, since all the past simulations performed with two flavors of Wilson fermions inside the Aoki phase were done with an external twisted mass term.
3. We could try to select one of these hypothetical new vacua by adding an external source proportional to $i\bar{\psi}\gamma_5\psi$, but this introduces a severe sign problem in the simulations.

The Aoki phase without external sources is very hard to simulate. Inside the Aoki phase there are exactly massless pions and quasi-zero modes, rendering the condition number of the Wilson Dirac operator quite high. A standard solver will not be enough to invert the Dirac operator in a HMC simulation. Fortunately, in recent years a number of efficient solvers have appeared, and by using a DD-HMC [38] we could invert the Dirac operator at a reasonable speed with the resources

available to us, i.e., the clusters of the department of theoretical physics of the University of Zaragoza, comprising 160 cores and 280 Gb of available memory, interconnected by a gigabit network. The simulations were parallelized for 4, 8 or 12 cores using openMP, and we did several simultaneous runs. The iteration count of the solver ranged from a handful outside the Aoki phase, to a few hundred inside the Aoki phase in the largest volume 8^4 and without external sources. For volumes higher than 8^4 , the inversion time became prohibitive for our computing resources and our time-frame, and we should look for new ways of simulating the Aoki phase.

Moreover, there is an additional problem with these simulations: since quasi-zero modes appear inside the Aoki phase, the eigenvalues sometimes attempt to cross the origin, and they would do so, were it not for the fact that the crossing of eigenvalues is forbidden by the HMC dynamics: the potential energy in the molecular dynamics step diverges at the origin, and there is an infinite repulsion that prevents the eigenvalues from crossing.

In order to solve this problem we classified our simulations according to its sector number, i.e., the number of 'crossed' eigenvalues of the hermitian Dirac-Wilson operator they had: beginning from a completely symmetric state (same number of positive and negative eigenvalues), the number of the eigenvalues that should cross to reach the desired state; and we performed simulations with a different number of crossed eigenvalues for each volume. Now this does not completely solve the problem, since we still don't know the weight of each sector within the partition function. The only way to simulate dynamically all the sectors is to add a twisted mass term, but as explained above we are primarily interested in the results without external sources. Fortunately the weights of the different sectors evolve very slowly with the twisted mass, and we can then extrapolate the data to vanishing twisted mass.

Indeed in Table 1 we report the weights of the different sectors as a function of the volume and twisted mass m_t . The values marked with * were obtained from reweighting and extrapolation² of the configurations generated at $m_t = 0.01$ in the case of the 4^4 , and $m_t = 0.05$ in the case of the 6^4 (the $m_t = 0.01$ simulation for the 6^4 came later, and has less statistics), *Qch* stands for quenched, whereas the numbers marked as MFA come from a *MFA* (Microcanonical Fermion Average) inspired approach [39], and were obtained by diagonalizing 4×10^6 gauge configurations. As can be seen in this table the weights show a very mild m_t -dependence, although the 4^4 results at $m_t = 0$ obtained from *MFA* simulations are at two standard deviations from the extrapolated results. It is not clear to us if this discrepancy has a statistical origin or reflects a discontinuity of the weights of different sectors at $m_t = 0$. However if the latter were the actual case, the standard picture for the Aoki phase could not be realized since it requires continuity in the sample of relevant gauge configurations at $m_t = 0$. Hence we will assume continuity of the

²Both methods agreed for our data, except for the higher sectors (2,3) of the 6^4 , where the low statistics prevented reweighting from working properly

weights, and in the following sections we will use Table 1 to reconstruct the PDF for the interesting bilinears by summing the weighted PDF's of all sectors for a given volume. The details of the runs can be checked in Table 2.

Vol	m_t	Sector 0	Sector 1	Sector 2	Sector 3+
4^4	MFA	$80\% \pm 6\%$	$20\% \pm 6\%$	0%	0%
4^4	0.0*	$67.6\% \pm 1.3\%$	$32.2\% \pm 1.4\%$	$0.11\% \pm 0.07\%$	0%
4^4	0.01	$68.1\% \pm 1.0\%$	$31.6\% \pm 1.0\%$	$0.24\% \pm 0.08\%$	0%
4^4	0.05	$70.4\% \pm 0.3\%$	$29.4\% \pm 0.3\%$	$0.18\% \pm 0.07\%$	0%
4^4	0.10	$69.3\% \pm 0.9\%$	$30.4\% \pm 0.9\%$	$0.28\% \pm 0.04\%$	0%
4^4	0.30	$70.3\% \pm 0.7\%$	$29.6\% \pm 0.6\%$	$0.14\% \pm 0.05\%$	0%
4^4	0.50	$70.0\% \pm 0.5\%$	$29.8\% \pm 0.5\%$	$0.20\% \pm 0.06\%$	0%
4^4	Qch	$71.4\% \pm 0.3\%$	$28.5\% \pm 0.2\%$	$0.07\% \pm 0.02\%$	0%
6^4	0.0*	$36\% \pm 4\%$	$59.2\% \pm 2.3\%$	$4.77\% \pm 0.45\%$	$0.12\% \pm 0.05\%$
6^4	0.01	$51.8\% \pm 0.6\%$	$44.7\% \pm 0.7\%$	$3.3\% \pm 0.3\%$	0%
6^4	0.05	$49.9\% \pm 0.8\%$	$45.7\% \pm 0.4\%$	$4.3\% \pm 0.5\%$	$0.08\% \pm 0.02\%$
6^4	0.10	$51.4\% \pm 0.6\%$	$44.7\% \pm 0.6\%$	$3.86\% \pm 0.07\%$	$0.06\% \pm 0.02\%$
6^4	0.30	$52.7\% \pm 1.2\%$	$44.3\% \pm 1.0\%$	$2.96\% \pm 0.20\%$	$0.06\% \pm 0.06\%$
6^4	0.50	$53.1\% \pm 0.6\%$	$44.0\% \pm 0.5\%$	$2.82\% \pm 0.18\%$	$0.06\% \pm 0.02\%$
6^4	Qch	$55.0\% \pm 1.6\%$	$43.0\% \pm 1.9\%$	$2.0\% \pm 0.4\%$	0%
8^4	0.05	$36.0\% \pm 0.6\%$	$48.2\% \pm 0.7\%$	$13.9\% \pm 0.4\%$	$2.06\% \pm 0.20\%$
8^4	0.10	$36.1\% \pm 0.6\%$	$47.0\% \pm 0.4\%$	$14.6\% \pm 0.3\%$	$2.22\% \pm 0.28\%$
8^4	0.30	$38.2\% \pm 0.4\%$	$46.8\% \pm 0.4\%$	$13.33\% \pm 0.12\%$	$1.6\% \pm 0.4\%$
8^4	0.50	$39.0\% \pm 0.3\%$	$48.5\% \pm 0.3\%$	$11.33\% \pm 0.25\%$	$1.18\% \pm 0.11\%$

Table 1: Weights of the different sectors as a function of the volume and m_t . The values marked with * were obtained from reweighting and extrapolation of the configurations generated at $m_t = 0.01$ for the 4^4 and $m_t = 0.05$ for the 6^4 .

The high variability of the acceptance ratio is due to the hard task of fine-tuning the solver parameters and the trajectory length to work properly for each case. As the table shows, certain simulations required a very small time-step for the HMC to work; in those simulations the eigenvalues were trying to cross the origin, generating high forces in the molecular dynamics, and thence creating a high rejection rate, so this small time-step was strictly necessary to obtain reasonable data. To enhance acceptance, we introduced the *replay trick* in the hardest calculations, i.e., those inside the Aoki phase and without a twisted mass term. During replays, the stepsize was halved whereas the trajectory length was kept constant.

In order to calculate the PDF's we diagonalized the hermitian Dirac-Wilson operator on all configurations generated and found the eigenvalues. The diagonalizations were carried out by using a parallelized Lanczos algorithm, combined with a Sturm bisection. The algorithm was checked heavily against the LAPACK library before starting production to ensure that our results were correct, but our algorithm was much faster than the LAPACK standard diagonalization procedure for hermitian matrices, and used a small fraction of the memory, because in our algorithm the fermionic matrix was generated on-the-fly. The diagonalization times ranged from around 100 seconds for the 4^4 to a few hours (close to four) in the case

Vol	β	κ	m_t	Sector	Confs	Acc (%)	HMC Step	l_{Traj}
4^4	2.00	0.25	0.00	0	5000	99% (72%)	3.125e-03	0.1
4^4	2.00	0.25	0.00	1	2473	99% (73%)	2.5e-03	0.1
6^4	2.00	0.25	0.00	0	676	92% (60%)	6.25e-04	0.04
6^4	2.00	0.25	0.00	1	2000	88% (56%)	6.25e-04	0.04
4^4	4.00	0.18	0.00	<i>All*</i>	10000	92%	0.1	1.0
6^4	4.00	0.18	0.00	<i>All*</i>	5000	84%	0.1	1.0
8^4	4.00	0.18	0.00	<i>All*</i>	5000	74%	0.1	1.0
4^4	2.00	0.25	0.01	<i>All</i>	5001	83%	0.0133	0.2
4^4	2.00	0.25	0.05	<i>All</i>	5001	86%	0.05	0.4
4^4	2.00	0.25	0.10	<i>All</i>	5664	92%	0.025	0.5
4^4	2.00	0.25	0.30	<i>All</i>	4999	96%	0.1	0.5
4^4	2.00	0.25	0.50	<i>All</i>	10000	99%	0.025	0.5
6^4	2.00	0.25	0.01	<i>All</i>	2520	99% (89%)	0.01	0.4
6^4	2.00	0.25	0.05	<i>All</i>	5000	70%	0.05	0.4
6^4	2.00	0.25	0.10	<i>All</i>	5000	95%	0.05	0.4
6^4	2.00	0.25	0.30	<i>All</i>	5001	90%	0.1	0.5
6^4	2.00	0.25	0.50	<i>All</i>	5001	99%	0.05	0.4
8^4	2.00	0.25	0.05	<i>All</i>	5001	77%	0.04	0.2
8^4	2.00	0.25	0.10	<i>All</i>	5001	84%	0.05	0.5
8^4	2.00	0.25	0.30	<i>All</i>	5001	96%	0.05	0.4
8^4	2.00	0.25	0.50	<i>All</i>	10000	98%	0.05	0.5

Table 2: Details of the simulations that generated our configurations. The word *Sector* refers to the eigenvalue sector where the simulation was performed, as explained above. The * means that only sector 0 contributes in this case, so all sectors were taken into account. Also, *Confs* refers to the number of configurations saved (we saved one configuration for each two generated, so as to reduce the autocorrelations), *Acc* is the acceptance ratio of the simulation (the value in parenthesis marks the acceptance without the replay trick), *HMC Step* is the molecular dynamics timestep used (during replays, the step was halved) and l_{Traj} is the trajectory length used.

of the 8^4 on our 12 core machines.

4 Probability Distribution Function of Q

First of all, we checked the behavior of the PDF of the operator Q outside the Aoki phase (point at $\beta = 4.0$ and $\kappa = 0.18$) in the three volumes considered (4^4 , 6^4 and 8^4). As figure 2 clearly indicates, the operator Q behaves as an intensive operator, as expected, and its PDF tends to a Dirac delta at the origin as the volume increases, showing that both parity and P' symmetries are realized in the vacuum

of the physical phase.




Figure 2: PDF of operator Q outside the Aoki phase for three different volumes.

Let's see what happens inside the Aoki phase ($\beta = 2.0$ and $\kappa = 0.25$ and for this case we will use data from the 4^4 and 6^4 volumes (the 8^4 inside the Aoki phase was very expensive for us). We see in figure 3 and figure 4 how the PDF behaves quite differently depending on the volume and on the sector. However, when we compute the final result taking into account the weight of each sector, we see in figure 5 how all these different plots converge to a single peak of constant width, in remarkable contrast with the results in the physical phase reported in figure 2. Therefore we expect, according to our discussion in Section 2.4, that not only parity but also P' will be spontaneously broken inside the Aoki phase. We should notice that this result needs to hold if the standard picture of the Aoki phase is correct, since otherwise $\langle c_0^2 \rangle = \langle c_3^2 \rangle \neq 0$ in this phase.

Of course we are far from the thermodynamic limit, and one might argue that, as stated in Table 1, the contribution of sector 2 to the 6^4 volume is important enough to be taken into account. This sector was not simulated because it was extremely expensive from the numerical point of view, and since its weight is less than 5%, we don't expect any changes to be relevant to the final result.

Another indication that there are new vacua inside the Aoki phase not considered in the standard picture, comes from measuring the PDF of the operator Q on configurations dynamically generated with several values of an external twisted




Figure 3: PDF of operator Q inside the Aoki phase for the two sectors considered in the 4^4 case. The sector 1 clearly shows a double peaked behavior

mass term, $m_t i\bar{\psi}\gamma_5\tau_3\psi$, but keeping $m_t = 0$ in the definition of Q . Since the addition of an external source selects a standard Aoki vacuum, we expect that all the contributions to the PDF coming from the other vacua will be removed. As we can see in figure 6, the PDF for this case is essentially different than the one shown in figure 5.

First of all, the PDF seems to be independent of the value of the external field. We might appreciate a slight dependence on the volume, since it seems that the peak height decreases as V increases, but this effect is too small to be significant, and in any case it is a good approximation to say that the distribution is also independent of the volume. Comparing this distribution with the former one (Gibbs state, without external source, figure 5), we notice that the PDF of the operator with external source is much wider, and we then expect the spectrum to be different as well. The essential difference lies in the low modes of the Dirac operator: in the case without external source, there is a lower bound given by $1/V$ for the modulus of any eigenvalue, whereas at $m_t \neq 0$ the eigenvalues can become arbitrarily small³. Since the PDF depends strongly on the spectrum of the Dirac operator, we

³A consequence of this property is the fact that the PDF of Q at $m_t \neq 0$ fits perfectly to a Lorentzian distribution of infinite tails. One could also be concerned that Q become singular if a




Figure 4: PDF of operator Q inside the Aoki phase for the two sectors considered, now in the 6^4 volume. The double peak of sector 1 becomes narrower, but we should expect wider contributions from higher sectors.

expect the cases $m_t = 0$ and $m_t \neq 0$ to be inherently different.

We think that this result is the strongest indication of additional vacua structure in the Aoki phase outside the standard picture. Indeed the results of figure 7 clearly show that the sample of gauge configurations obtained in the Aoki phase at $m_t = 0$ is qualitatively different from the sample obtained at $m_t \neq 0$, even in the $m_t \rightarrow 0$ limit. On the other hand the differences in the samples can never come from the additional standard Aoki vacua since if we change the twisted mass term in the dynamical generation by any other symmetry breaking term selecting other standard Aoki vacuum, for instance $im_t \bar{\psi} \gamma_5 \tau_{1,2} \psi$, the fermion determinant does not change.

5 Second moments of the PDFs of fermion bilinears

Since P' is broken according to our results plus the assumption that Q is an intensive operator, the symmetry arguments supporting the existence of a tower of

zero-mode appeared in the spectrum. Actually, the set of configurations with a zero-mode has measure zero, and Q remains non-singular.




Figure 5: PDF of operator Q inside the Aoki phase for two different volumes. The differences in the area can be explained by realizing that sector 2 is missing in the 6^4 volume, which should account for a 5% of the area.

sum rules are no longer valid, thus it is natural to explore if the expectation value $\langle (i\bar{\psi}\gamma_5\psi)^2 \rangle$ will be non-zero in the Gibbs state inside the Aoki phase. This observable is extremely difficult to measure, nonetheless we managed to obtain sensible results from our data, which we show in Table 3.

The fourth column in this table refers to $\langle (i\bar{\psi}\gamma_5\tau_3\psi)^2 \rangle$, the landmark of the Aoki phase. As we see, it is clearly non-zero for all the volumes, confirming that our simulations lie within the Aoki phase. The second column shows the values for $\langle (i\bar{\psi}_u\gamma_5\psi_u)^2 \rangle$, which is an order parameter for parity, but it takes into account just one flavor (which we labeled u). This quantity should be non-zero inside the parity-breaking Aoki phase, regardless of the discussion of the new vacua. Finally, the most important observable, the flavor singlet pseudoscalar $\langle (i\bar{\psi}\gamma_5\psi)^2 \rangle$, which – since we are dealing with two degenerate flavors, u and d – is the sum of two of the former condensates $i\bar{\psi}_u\gamma_5\psi_u + i\bar{\psi}_d\gamma_5\psi_d$. The standard picture of the Aoki phase predicts zero expectation value of this parameter in any Aoki vacuum, whereas each one of the pseudoscalars restricted to one flavor $i\bar{\psi}_{u,d}\gamma_5\psi_{u,d}$ will be non zero due to parity breaking. Hence, the standard picture of the Aoki phase enforces




Figure 6: PDF of operator Q inside the Aoki phase for configurations generated with a twisted mass source.

an antiferromagnetic ordering of the pseudoscalars $i\bar{\psi}_u\gamma_5\psi_u = -i\bar{\psi}_d\gamma_5\psi_d$, which is not required in our hypothesis of the new vacua. The data of the third column in Table 3 show a clear non-zero expectation value in the case of the largest volume 6^4 , supporting our previous discussion regarding P' breaking.

Concerning equation (22), which relates the mean value of the square Aoki condensate in the Gibbs state (ε -regime) with the expectation value of this order parameter in the vacuum selected by a twisted-mass term (p -regime), and since we have data for the Aoki condensate at several lattice sizes and twisted masses (see Table 4), we can check its plausibility. This is a relevant test since, as discussed in Section 2.3, equation (22) should be realized in the standard Aoki scenario [22, 27]. To this end we report in figure 8 our results for the Aoki condensate at several values of the twisted-mass, m_t , in 4^4 , 6^4 and 8^4 lattices. The circles in the ordinates axis stand for the values of the Aoki condensate at $m_t = 0$ in 4^4 and 6^4 lattices, obtained from equation (22), and using as input our results for $\langle(i\bar{\psi}\gamma_5\tau_3\bar{\psi})^2\rangle$ reported in Table 3. The figure, which is a Fisher plot [40], shows that any reliable extrapolation of the data to $m_t = 0$ gives a value for the Aoki condensate larger than the one obtained from equation (22).

One could argue that, at the volumes we are working, the finite size effects should be large and noticeable, and that the inclusion of these effects could lead to

Volume	$\langle (i\bar{\Psi}_u \gamma_5 \Psi_u)^2 \rangle$	$\langle (i\bar{\Psi} \gamma_5 \Psi)^2 \rangle$	$\langle (i\bar{\Psi} \gamma_5 \tau_3 \Psi)^2 \rangle$	$\langle \mathcal{Q}^2 \rangle$
4 ⁴ Sec 0	$(1.93 \pm 0.02) 10^{-2}$	$(2.51 \pm 0.07) 10^{-2}$	$(5.23 \pm 0.09) 10^{-2}$	$(6.8 \pm 0.3) 10^{-3}$
6 ⁴ Sec 0	$(2.15 \pm 0.05) 10^{-2}$	$(2.82 \pm 0.18) 10^{-2}$	$(5.76 \pm 0.13) 10^{-2}$	$(7.4 \pm 0.6) 10^{-3}$
4 ⁴ Sec 1	$(6.0 \pm 0.5) 10^{-3}$	$(-5.5 \pm 0.5) 10^{-2}$	$(7.9 \pm 0.4) 10^{-2}$	$(3.34 \pm 0.20) 10^{-2}$
6 ⁴ Sec 1	$(1.75 \pm 0.09) 10^{-2}$	$(0.4 \pm 0.4) 10^{-2}$	$(6.56 \pm 0.26) 10^{-2}$	$(1.53 \pm 0.15) 10^{-2}$
4 ⁴ Total	$(1.50 \pm 0.03) 10^{-2}$	$(-0.6 \pm 1.9) 10^{-3}$	$(6.08 \pm 0.19) 10^{-2}$	$(1.54 \pm 0.08) 10^{-3}$
6 ⁴ Total	$(1.81 \pm 0.12) 10^{-2}$	$(1.27 \pm 0.29) 10^{-2}$	$(6.0 \pm 0.3) 10^{-2}$	$(1.17 \pm 0.13) 10^{-3}$

Table 3: Interesting v.e.v. for the Aoki phase in the different sectors. Only statistical errors are shown.

V	m_t	$\langle i\bar{\Psi} \gamma_5 \tau_3 \Psi \rangle$
4 ⁴	0.01	$(56.00 \pm 0.06) 10^{-2}$
4 ⁴	0.05	$(65.149 \pm 0.022) 10^{-2}$
6 ⁴	0.01	$(53.603 \pm 0.015) 10^{-2}$
6 ⁴	0.05	$(66.888 \pm 0.008) 10^{-2}$
8 ⁴	0.05	$(66.953 \pm 0.007) 10^{-2}$
4 ⁴	0.10	$(75.512 \pm 0.009) 10^{-2}$
6 ⁴	0.10	$(76.291 \pm 0.004) 10^{-2}$
8 ⁴	0.10	$(76.307 \pm 0.003) 10^{-2}$
4 ⁴	0.30	$(92.214 \pm 0.005) 10^{-2}$
6 ⁴	0.30	$(92.309 \pm 0.002) 10^{-2}$
8 ⁴	0.30	$(92.311 \pm 0.002) 10^{-2}$
4 ⁴	0.50	$(96.225 \pm 0.003) 10^{-2}$
6 ⁴	0.50	$(96.241 \pm 0.001) 10^{-2}$
8 ⁴	0.50	$(96.243 \pm 0.001) 10^{-2}$

Table 4: Evolution of the v.e.v. of the Aoki parameter as a function of the twisted mass external field.




Figure 7: PDF of operator Q inside the Aoki phase for all configurations. Comparison of the case $m_t \neq 0$ versus the case $m_t = 0$.

an agreement between our measurements at $m_t = 0$ and $m_t \neq 0$. Nonetheless, the data at $m_t = 0$ —which we expect to suffer more these finite volume effects due to the existence of massless pions—reveals that these effects are not very large, for the value of the Aoki condensate remains very stable after a fivefold increase in the volume.

6 Conclusions and Outlook

Three years ago the standard scenario for the phase structure of lattice QCD with two degenerate flavors of Wilson fermions was revised by three of us in [21], where we conjectured on the appearance of new vacua in the Aoki phase, which can be characterized by a non-vanishing vacuum expectation value of $i\bar{\psi}_u\gamma_5\psi_u + i\bar{\psi}_d\gamma_5\psi_d$, and which can not be connected with the Aoki vacua by parity-flavor symmetry transformations. However, Sharpe pointed out in [22] that the standard picture for the Aoki phase can be understood using the chiral effective theory appropriate to the Symanzik effective action, and that within this standard analysis, the flavor-singlet pseudoscalar expectation value vanishes, $\langle i\bar{\psi}\gamma_5\psi \rangle = 0$. As the standard scenario for the Aoki phase is indeed a direct consequence of the χPT application




Figure 8: Fisher plot of the evolution of the v.e.v. of the Aoki parameter as a function of the twisted mass external field. The circles show the prediction obtained from the simulations in the Gibbs state (see eq.(22)). The fit shown is meant to guide the eye, and shouldn't be taken as a prediction of the behavior of the v.e.v. of the Aoki parameter. The fit should be more reliable for small values of m_t .

to this phase, we were also calling into question in [21] the validity of the χ_{PT} analysis, at least for low values of β , and therefore large values of the lattice spacing a . These issues are relevant enough to make it worthwhile to continue these investigations in order to clarify the actual scenario for the Aoki phase.

For the last few years we have performed an extensive research on the vacuum structure of the Aoki phase, in order to find out if our alternative vacuum structure, derived from a PDF analysis, was realized or not. These simulations, which have been mainly performed in the absence of any parity-flavor symmetry breaking external source, are plagued by technical difficulties which have been responsible for the slow progress in the field. Indeed the addition of a twisted mass external source, as it has been done in the past simulations of the Aoki phase, automatically selects the vacuum where the standard properties of the Aoki phase are verified. This point could explain why nobody ever saw a new phase like the one we are proposing, since all the past simulations performed with two flavors of Wilson fermions inside the Aoki phase were done within an external twisted mass term.

Notwithstanding these difficulties, we have provided in this work evidence

pointing to a more complex vacuum structure in the Aoki phase of two-flavor QCD, as conjectured in [21]. Indeed the results reported in Table 3, which were obtained under the assumption that the weights of the different sectors are continuous at $m_t = 0$, show how we were able to perform a direct measurement of $\langle (i\bar{\psi}\gamma_5\psi)^2 \rangle$ in the Aoki phase, which gave a non-zero value for this operator in the 6^4 lattice, of the same order of magnitude as $\langle (i\bar{\psi}_u\gamma_5\psi_u)^2 \rangle$, and $\langle (i\bar{\psi}\gamma_5\tau_3\psi)^2 \rangle$, the last two being non-vanishing in the Aoki phase in the standard scenario. Thus this result points to the breaking of the hypothesis of the sum-rules (13). Furthermore a check of equation (22), a equation which should hold if the standard scenario for the Aoki phase is realized, reported in figure 8, points out to a more complex vacuum structure too.

However the strongest indication, in our opinion, on the existence of a vacuum structure in the Aoki phase, more complex than the one of the standard picture, comes from our analysis of the PDF of the operator Q (23). Our motivation for the analysis of this kind of density of "topological charge" operator, which measures the asymmetries in the eigenvalue distribution of the hermitian Dirac-Wilson operator, was twofold. First it appears in the computation of the second moment of the PDF of the flavor singlet pseudoscalar order parameter (12), and second Q is an order parameter for the symmetry P' , composition of parity and discrete flavor transformations, described in Section 2.

Our numerical results for the PDF of Q , together the assumption that Q is an intensive operator, suggest that the P' symmetry is realized in the vacuum of the physical phase, but not in the Aoki phase, at least in the strong coupling region we have analyzed. However the cleanest signal of further structure in the Aoki phase comes from the results on the PDF of Q reported in figure 7. Those results clearly show that the sample of gauge configurations obtained in the Aoki phase at $m_t = 0$ is qualitatively different from the sample obtained at $m_t \neq 0$, since they give incompatible PDF's. Furthermore this result stays true in the $m_t \rightarrow 0$ limit, as follows from the independence of the shape of the PDF of Q on the twisted mass m_t . On the other hand the differences in the samples can never come from the additional standard Aoki vacua since if we change the twisted mass term in the dynamical generation by any other symmetry breaking term selecting other standard Aoki vacuum, as for instance $im_t\bar{\psi}\gamma_5\tau_{1,2}\psi$, the fermion determinant does not change.

As the sum rules (13) required in the standard scenario are only supported, as discussed before, by the χPT analysis of the Aoki phase, our results could call into question also the validity of the χPT analysis performed in [11] for low values of β (around 2.0, very coarse lattices). However, χPT is expected to work at higher values of β , near the continuum limit. Indeed, one could be concerned that, at such a low β , we are far from the continuum limit. Nonetheless our analysis is devoted to the Aoki phase on the lattice, which might not even have a continuum limit.

In any case any improvement of our results in larger lattices and for more β, κ values would be of great interest. Unfortunately, given the technical difficulties

of the simulations inside the Aoki phase, this calculation is outside our present computing resources.

Acknowledgments

It is a pleasure to thank Steve Sharpe for his invaluable help, interesting comments and fruitful discussions; his remarks allowed us to improved greatly this work. We also want to thank Fabrizio Palumbo for useful discussions. This work was funded by an INFN-MICINN collaboration (under grant AIC-D-2011-0663), MICINN (under grants FPA2009-09638 and FPA2008-10732), DGIID-DGA (grant 2007-E24/2) and by the EU under ITN-STRONGnet (PITN-GA-2009-238353). E. Follana is supported on the MICINN Ramón y Cajal program, and A. Vaquero is supported by the Research Promotion Foundation (RPF) of Cyprus under grant ΠΡΟΣΕΛΚΥΣΗ/ΝΕΟΣ/0609/16.

References

- [1] V. Azcoiti, A. Nakamura, Phys. Rev. **D27**, 2559 (1983).
- [2] H.W. Hamber, Nucl. Phys. **B251**, 182 (1985).
- [3] S. Aoki, Phys. Rev. **D30**, 2653 (1984).
- [4] S. Aoki, Phys. Rev. Lett. **57**, 3136 (1986).
- [5] S. Aoki, S. Boettcher, and A. Gocksch, BNL-ABG-1; hep-lat/9312084 (1993).
- [6] M. Creutz, arXiv:hep-lat/9608024.
- [7] K.M. Bitar, Phys. Rev. **D56**, 2736 (1997).
- [8] S. Aoki, T. Kaneda, and A. Ukawa, Phys. Rev. **D56**, 1808 (1997).
- [9] K.M. Bitar, U.M. Heller, and R. Narayanan, Phys. Lett. **B418**, 167 (1998).
- [10] R.G. Edwards, U.M. Heller, R. Narayanan, and R.L. Singleton, Jr, Nucl. Phys. **B518**, 319 (1998).
- [11] S.R. Sharpe, R.L. Singleton, Jr, Phys. Rev. **D58**, 074501 (1998).
- [12] S. Aoki, Nucl. Phys. **B** (Proc. Suppl.) **60A**, 206 (1998).
- [13] K. Bitar, Nucl. Phys. **B** (Proc. Suppl.) **63A-C**, 829 (1998).
- [14] S. Sharpe, R.L. Singleton Jr, Nucl. Phys. **B** (Proc. Suppl) **73**, 234 (1999).
- [15] R. Kenna, C. Pinto, and J.C. Sexton, Phys. Lett. **B505**, 125 (2001).

- [16] M. Golterman, Y. Shamir, Phys. Rev. **D68**, 074501 (2003).
- [17] E.M. Ilgenfritz, W. Kerler, M. Müller-Preussker, A. Sternbeck, and H. Stuben, Phys. Rev. **D69**, 074511 (2004).
- [18] A. Sternbeck, E.M. Ilgenfritz, W. Kerler, M. Müller-Preussker, and H. Stuben, Nucl. Phys. **B** (Proc. Suppl.) **129&130**, 898 (2004).
- [19] M. Golterman, S.R. Sharpe, and R.L. Singleton, Jr, Nucl. Phys. **B** (Proc. Suppl.) **140**, 335 (2005).
- [20] V.E.M. Ilgenfritz, M. Müller-Preussker, M. Petschlies, K. Jansen, M.P. Lombardo, O. Philipsen, L. Zeidlewicz, and A. Sternbeck, Proc. Sci., Lattice2007, (2008) 323.
- [21] V. Azcoiti, G. Di Carlo, A. Vaquero, Phys. Rev. **D79**, 014509 (2009).
- [22] S. Sharpe, Phys. Rev. **D79**, 054503 (2009)
- [23] A. Vaquero, V. Azcoiti, G. Di Carlo, E. Follana, PoS **LATTICE2009** 068 (2009)
- [24] V. Azcoiti, G. Di Carlo, E. Follana, A. Vaquero, PoS **LATTICE2010** 091 (2010)
- [25] V. Azcoiti, G. Di Carlo, E. Follana, A. Vaquero, PoS **LATTICE2011** 112 (2011)
- [26] G. Akemann, P.H. Damgaard, K. Splittorff, J.J.M. Verbaarschot Phys. Rev. **D83**, 085014 (2011)
- [27] V. Azcoiti, V. Laliena, and X.Q. Luo, Phys. Lett. **B354**, 111 (1995).
- [28] V. Azcoiti, G. Di Carlo, and A. Vaquero, JHEP**0804:035**, (2008).
- [29] T. Banks, A. Casher, Nucl. Phys. **B169**, 103 (1980).
- [30] H. Leutwyler, A. Smilga, Phys. Rev. D **46**, 5607 (1992).
- [31] J. Smit, J.C. Vink, Nucl. Phys. **B286**, 485 (1987).
- [32] P. Hasenfratz, V. Laliena, F. Niedermayer, Phys. Lett. **B427**, 125 (1998).
- [33] C. Vafa, E. Witten, Phys. Rev. Lett. **53**, 535 (1984).
- [34] V. Azcoiti, A. Galante, Phys. Rev. Lett. **83**, 1518 (1999).
- [35] X. Ji, Phys. Lett. **B554**, 33 (2003).
- [36] P.R. Crompton, Phys. Rev. D **72**, 076003 (2005).

- [37] V. Azcoiti, G. Di Carlo, E. Follana, A. Vaquero, JHEP**1007:047**, (2010).
- [38] M. Luscher, Comput. Phys. Commun. **156**, (2004) 20.
- [39] V. Azcoiti, G. Di Carlo, A.F. Grillo, Phys. Rev. Lett. **65**, 2239 (1990).
- [40] J.S. Kouvel, M.E. Fisher, Phys. Rev. **136**, A1626 (1964).